

UNFOLDING OF CHAOTIC QUADRATIC MAPS — PARAMETER DEPENDENCE OF NATURAL MEASURES

HANS THUNBERG

ABSTRACT. We consider perturbations of quadratic maps f_a admitting an absolutely continuous invariant probability measure, where a is in a certain positive measure set $\mathcal{A}$ of parameters, and show that in any neighborhood of any such an f_a , we find a rich fauna of dynamics. There are maps with periodic attractors as well as non-periodic maps whose critical orbit is absorbed by the continuation of any prescribed hyperbolic repeller of f_a . In particular, Misiurewicz maps are dense in $\mathcal{A}$.

Almost all maps f_a in the quadratic family is known to possess a unique natural measure, that is, an invariant probability measure μ_a describing the asymptotic distribution of almost all orbits. We discuss weak*-(dis)continuity properties of the map $a \mapsto \mu_a$ near the set $\mathcal{A}$, and prove that almost all maps in $\mathcal{A}$ have the property that μ_a can be approximated with measures supported on periodic attractors of certain nearby maps. On the other hand, for any $a \in \mathcal{A}$ and any periodic repeller Γ_a of f_a , the singular measure supported on Γ_a can also be approximated with measures supported on nearby periodic attractors. It follows that $a \mapsto \mu_a$ is not weak*continuous on any full-measure subset of $(0, 2]$. Some of these results extend to unimodal families with critical point of higher order, and even to not-too-flat flat topped families.

1. DEFINITIONS, STATEMENTS AND RELATED RESULTS

1.1. Introduction. An invariant measure μ for an interval map f is called a *natural (physical, Sinai-Ruelle-Bowen)* measure, if μ describes the asymptotic distribution of $\{f^n(x)\}_{n=0}^\infty$ for all x in a set of positive Lebesgue measure, i.e. if

$$\mu \stackrel{\text{weak}^*}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \delta_{f^k(x)}$$

1991 *Mathematics Subject Classification.* Primary: 58F03, 58F11; Secondary: 26A18.

Key words and phrases. one-dimensional dynamics, unimodal families, unfolding of chaotic maps, parameter dependence of natural measures.

holds for all x in a positive measure set. Those x for which this do hold, are said to be *generic* for μ .

We consider the quadratic family $f_a(x) = 1 - ax^2$, $0 < a \leq 2$, of dynamical systems on $I = [-1, 1]$. If f_a admits an *acip* (absolutely continuous invariant probability measure) μ_a , then μ_a is a natural measure for f_a , describing the asymptotics of almost all orbits (Theorem V.1.5, [MS93]). If f_a has a periodic attractor $\{x_i\}_{i=1}^p$, then

$$\mu_a := \frac{1}{p} \sum_{i=1}^p \delta_{x_i}$$

is again a natural measure for f_a which describes the asymptotics for almost all points ([BL91]).

For a long time it was conjectured that for almost all a , f_a either has a periodic attractor or admits an acip. This fundamental result was recently proved by M. Lyubich ([Lyu97]). It follows that the mapping

$$\Psi : a \mapsto \mu_a =: \text{the natural measure of } f_a$$

is well defined for almost all a in $(0, 2]$. Of course, Ψ is continuous on the set of hyperbolic attracting maps. In what follows we discuss the structure in parameter space and the parameter dependence of μ_a near certain maps admitting an acip, more precisely the maps f_a considered by Benedicks and Carleson.

Theorem (Benedicks and Carleson, [BC85], [BC91]). *There exists a set $\mathcal{A} \subset (0, 2]$ of positive Lebesgue measure, with 2 a Lebesgue density point of $\mathcal{A}$, such that if $a \in \mathcal{A}$, then the Lyapunov exponent of f_a at the critical value is positive and also f_a admits an acip μ_a with a density that belongs to L^p for any $1 \leq p < 2$. Furthermore, for almost all $a \in \mathcal{A}$, the critical point $c = 0$ is generic for μ_a .*

Jakobson, in [Jak81], was the first to prove the existence of a positive measure set of parameters admitting acips. Various versions and generalizations can be found in [Ryc88], [BY92], [TTY92], [Tsu93b], [Tsu93a], [MS93], [Thu97] and [Luz98].

1.2. Theorems and corollaries.

Theorem A. *For each $a \in \mathcal{A}$, there exists a sequence $\{a_n\}_{n=1}^\infty$ such that f_{a_n} has a super-stable periodic attractor of length r_n , such that*

- (i) $a_n \rightarrow a$, $n \rightarrow \infty$;
- (ii) $r_n \uparrow \infty$, $n \rightarrow \infty$;
- (iii) *if $x = 0$ is generic for μ_a , then $\mu_{a_n} \xrightarrow{\text{weak}^*} \mu_a$, $n \rightarrow \infty$.*

In [Thu96] a version of Theorem A is proven, assuming only sub-exponential growth of the derivative along the orbit of the critical value. In [Ure95] and [Ure96], Ures proves corresponding statements for the Hénon family.

Theorem B. *Let $\Gamma = \Gamma_a$ be a hyperbolic set for f_a , $a \in \mathcal{A}$, let $z = z(a)$ be any point in Γ and let Γ_b and $z(b)$ be the continuations of Γ and z . Then*

$$a \in \text{cl}\{b \mid f_b^N(0) = z(b) \text{ for some } N = N(b)\}$$

Remark 1. An obvious choice is $\Gamma_a = \{z(a)\}$, where $z(a)$ is the interior unstable fixed point. This implies that (this type of) Misiurewicz points are dense in $\mathcal{A}$. By taking $\mathcal{A}$ sufficiently close to 2, the theorem holds for any hyperbolic set of f_2 .

Theorem C. *Let $\{x_1, x_2, \dots, x_p\}$ be a hyperbolic periodic repeller for f_a , $a \in \mathcal{A}$. Then there exists a sequence of parameters $\{a_n\}_{n=1}^\infty$ converging to a such that f_{a_n} has a super-attractor and such that*

$$\mu_{a_n} \xrightarrow{\text{weak}^*} \mu_a^{\text{sing}} =: \frac{1}{p} \sum_{i=1}^p \delta_{x_i}.$$

We also state a theorem that holds for any post-critically finite Misiurewicz map in the quadratic family.

Theorem D. *Suppose f_a is a quadratic map whose critical orbit is pre-periodic to an unstable periodic orbit $\{x_1, x_2, \dots, x_p\}$. Then there is a sequence of parameters $\{a_n\}_{n=1}^\infty$ accumulating at a such that f_{a_n} has a super-stable period attractor and*

$$\mu_{a_n} \xrightarrow{\text{weak}^*} \mu_a^{\text{sing}} =: \frac{1}{p} \sum_{i=1}^p \delta_{x_i}.$$

Remark 2. Since $\Psi : a \mapsto \mu_a$ is continuous when restricted to a periodic window, the statements of theorems A, C and D remains true for any sequence of parameters running thru the periodic windows $J_n \ni a_n$.

From Theorem C and Remark 2, we obtain

Corollary 1. *$\Psi : a \mapsto \mu_a$ is not continuous at any point in $\mathcal{A}$, and Ψ is not continuous on any full-measure subset of $(0, 2]$.*

Since Misiurewicz maps admits an acip under some mild conditions, fulfilled in the quadratic family, [Mis81] and [BM89], Theorem D gives the following:

Corollary 2. *$\Psi : a \mapsto \mu_a$ is discontinuous at every post-critically finite Misiurewicz map in the quadratic family.*

Most of this can be generalized to generic unimodal families with critical point of higher order, and even to those flat-topped families that are considered in [Thu97]. What may happen is that the last statement of Theorem A becomes empty; it is not known if the critical point is generic for a positive set of acip-maps in these cases.

1.3. Related results. The measures μ_a , $a \in \mathcal{A}$, are known to be stable under random perturbations of the iterations of f_a ([BY92], [BV96]). Noise is thus in this way a savior, when modeling/experimenting with unimodal maps in the chaotic regime.

In [Ryc88], Rychlik gave a new proof of Jakobsons Theorem. A positive measure set $\mathcal{A}'$ of parameters is constructed such that for $a \in \mathcal{A}'$, f_a admits an acip μ_a with a density $\nu_a \in L^p$ for $1 \leq p < 2$. Then in [RS92], Rychlik and Sorret among other things proved that $a \mapsto \nu_a$, defined on $\mathcal{A}'$, is continuous in L^p , $1 \leq p < 2$, with a Hölder estimate at Misiurewicz points. The set $\mathcal{A}'$ also has full density at $a = 2$, and so it has a fat intersection with the set $\mathcal{A}$ considered in this paper.

As mentioned, Tsujii has a proof of the Benedicks-Carleson-Jakobson Theorem (leading to an a priori different positive measure set $\mathcal{A}''$ of Collet-Eckmann maps), and some generalizations there of, [Tsu93b] and [Tsu93a]. In [Tsu96] he also discusses weak*-continuity properties of the invariant measures, and among other things proves continuity of $\Psi|_{\mathcal{A}''}$ at Misiurewicz points. Also in this case, 2 is a Lebesgue density point of the good set $\mathcal{A}''$. He also constructs a set F of hyperbolic attracting maps accumulating on $a = 2$, whose natural measures converge to a point mass at the unstable fixed point at $x = -1$ as a tends to 2, and shows that $|F \cap [2 - \epsilon, 2]| \gtrsim \epsilon^2$.

We also remark that the special role of the parameter $a = 2$ is not that special, similar statements holds near any post-critically finite Misiurewicz parameter.

The following is also relevant to this discussion, even though it is a measure 0 phenomena: In [HK90] one constructs

- an uncountable set of parameters, accumulating on $a = 2$, such that the corresponding maps do not have any natural measure at all;
- an uncountable set of parameters, accumulating on $a = 2$, such that the corresponding maps have natural measures $\mu_a = \delta_{z(a)}$.

The first examples of maps with no natural measures was given in [Joh87].

2. PROPERTIES OF THE SET $\mathcal{A}$

In sections 2.1 – 2.3 we recall some basic facts and definitions from the construction of the set $\mathcal{A}$ that will be used in the proofs. We just state the results that we need in the sequel, and statements do not appear in their logical order. For proofs and more details we refer to [BC85], [BC91] and the expositions in [MS93] and [Luz98]. Similar statements with similar proofs can also be found in [Thu97], dealing with a Benedicks-Carleson theorem for certain flat-top families.

2.1. A partition on the interval. A small neighborhood $I^* = (-\delta, \delta)$ of the critical point $x = 0$ is chosen. I^* is partitioned into subintervals

$$I^* \setminus \{0\} = \bigcup_{|\mu| \geq -\log \delta} I_\mu = \bigcup_{\substack{|\mu| \geq -\log \delta \\ 1 \leq \nu \leq \mu^2}} I_{\mu\nu},$$

where $I_\mu = [e^{-(\mu+1)}, e^{-\mu})$ for $\mu > 0$, $I_\mu = -I_{-\mu}$ for $\mu < 0$, and

$$I_\mu = \bigcup_{1 \leq \nu \leq \mu^2} I_{\mu\nu}$$

is a subdivision of I_μ into μ^2 intervals of equal length. $I_{\mu\nu}^+$ denotes the union of $I_{\mu\nu}$ and its two nearest neighbors. We also define $\hat{I}_\mu = (-e^{-\mu}, e^{-\mu})$.

2.2. Partitions in parameter-space and the mappings ξ_n . The set $\mathcal{A}$ is given as

$$\mathcal{A} =: \bigcap_{n \geq 0} A_n,$$

where A_n is a decreasing sequence of sets in a small one-sided neighborhood of $a = 2$. A_n is constructed by deleting from A_{n-1} according to two principles, which together guarantees that for $a \in A_n$,

$$(2.1) \quad |Df_a^j(f_a(0))| \geq e^{\lambda j}, \quad \forall j \leq n,$$

for some $\lambda > 0$ independent of n and a . One of the exclusion principles simply requires that

$$(2.2) \quad |f_a^n(0)| \geq e^{-\alpha n}.$$

for some suitable, small $\alpha > 0$.

The mappings ξ_n from parameter space to dynamical space are defined via

$$\xi_n(a) := f_a^n(0).$$

These mappings are expanding as long as f_a^n is, in the following sense:

Lemma 1. *There is a constant C such that for all a sufficiently close to 2 the following holds: If $|D_x f_a^j(1)| \geq e^{\lambda j}$ for all $j \leq k$ then*

$$\frac{1}{C} \leq \left| \frac{D_a \xi_{k+1}(a)}{D_x f_a^k(1)} \right| \leq C.$$

On each A_n there is a partition $\mathcal{P}_n$ into intervals; each $a \in \mathcal{A}$ is given as $a = \bigcap_{n \geq 0} \omega_n(a)$, where $\omega_n(a)$ is the element of $\mathcal{P}_n$ containing a . For each a there is a sequence of times $n_k(a)$, the *essential free return times*, with the following properties:

- $n_k \leq n < n_{k+1} \implies \omega_n = \omega_{n_k}$;
- $n \leq n_k, \xi_n(\omega_{n_k}) \cap I^* \neq \emptyset \implies \xi_n(\omega_{n_k}) \subset \text{some } I_{\mu\nu}^+$;
- $I_{\mu\nu} \subset \xi_{n_k}(\omega_{n_k}) \subset I_{\mu\nu}^+$, for some $I_{\mu\nu}$.

A free return is followed by a so called *bound period*, when $\xi_{n_k+j}(a) = f_a^{n_k+j}(0)$ shadows an initial segment $f_a^j(0)$ of the critical orbit closely. More precisely: If $I_{\mu\nu} \subset \xi_{n_k}(\omega_{n_k}) \subset I_{\mu\nu}^+$, then $\xi_{n_k+j}(\omega)$ is in a bound period as long as

$$(2.3) \quad \left| \bigcup_{a \in \omega} f_a^j(-e^{-\mu}, e^{-\mu}) \right| \leq e^{-2\alpha j}.$$

We use $p = p(\mu, \omega)$ to denote the length of a bound period.

As long as orbiting outside I^* , $\xi_n(\omega)$ grows exponentially (Lemma 3). The small derivative picked up at a return to I^* is compensated for during the bound period by an inductive argument, the net effect is in fact a weaker exponential growth. In particular bound periods are always of finite length, (Lemma 2). Let p_k temporally denote the length of the bound period following a return at time n_k . By definition, $n_{k+1}(a)$ is the smallest integer $j \geq n_k(a) + p_k(a)$ such that $\xi_j(\omega_{n_k}(a)) \supset \text{some } I_{\mu\nu} \subset I^*$. A return to I^* at some time j , $n_k + p_k \leq j < n_{k+1}$, when no $I_{\mu\nu}$ is covered, is called an *inessential free return*. Such returns are also followed by a bound period, after which comes a free period terminating in a new free return. It can be shown that an essential free return always occur after finitely many steps. The “dynamics” of $\{\xi_n\}$ is described more precisely in the following lemmas.

Lemma 2. *Suppose inequality (2.1) and condition (2.2) hold for all $j \leq n$ and all $a \in \omega$, and suppose that $\xi_n(\omega) \subset I_{\mu\nu}^+$ where $|\mu| \geq -\log \delta$ and δ is sufficiently small. Then there exists positive constants C_0 and C , independent of δ , such that*

- (i) *for all $a \in \omega$, all $y \in f_a(\hat{I}_\mu)$ and all $j \leq p$,*

$$\frac{1}{C_0} \leq \frac{|D_x f_a^j(y)|}{|D_x f_a^j(1)|} \leq C_0;$$

- (ii) $C|\mu| \leq p(\mu, \omega) \leq 3|\mu|/\lambda \leq 3\alpha n/\lambda < n/100$;
- (iii) $|(f_a^p)'(x)| \geq Ce^{\lambda p/4}, \quad \forall x \in I_\mu$;
- (iv) $|\xi_{n+p(\mu, \omega)}(\omega)| \geq Ce^{\lambda p/4} |\xi_n(\omega)|$;

Lemma 3. *There is a $\lambda_0 > 0$ such that for any small $\delta > 0$ and a_0 sufficiently close to 2 the following holds: Suppose that $\omega \subset [a_0, 2]$ is such that $\xi_{\hat{n}}(\omega) \subset I_{\mu\nu}$ and $\xi_n(\omega)$ are two consecutive free returns with return times $\hat{n}$ and n , $\hat{n} < n$. Also assume that $|Df_a^j(1)| \geq e^{\lambda j}$ for all $j < n$ and all $a \in \omega$. Then there is a constant C , independent of δ , such that the following holds:*

- (i) $|\xi_{n-k}(\omega)| \leq Ce^{-\lambda_0 k} |\xi_n(\omega)|, \quad \forall 1 \leq k \leq n - \hat{n} - p(\mu, \omega)$;
- (ii) $|\xi_n(\omega)| \geq 2 |\xi_{\hat{n}}(\omega)|$.

Furthermore there is a positive integer $N_0(\delta)$ such that for any ω close to 2,

- (iii) $\xi_{k+j}(\omega) \cap I^* = \emptyset, \quad j = 0, 1, \dots, N_0 \implies$
 $|\xi_{k+N_0}(\omega)| \geq e^{\frac{\lambda_0}{2} N_0} |\xi_k(\omega)|.$

Let $\text{HD-dist}(J, K)$ denote the Hausdorff distance between the sets J and K .

Lemma 4. *Suppose inequality (2.1) and condition (2.2) hold for all $j \leq n$ and all $a \in \omega$, and suppose that $\xi_n(\omega) \subset I_\mu$ where $|\mu| \geq -\log \delta$. If ω is sufficiently close to 2, then for each $a, b \in \omega$ we have*

- (i) $\text{HD-dist} \left(f_a^j \left(\hat{I}_\mu \right), f_b^j \left(\hat{I}_\mu \right) \right) < \frac{1}{1000} \left| f_a^j \left(\hat{I}_\mu \right) \right|$
- (ii) $\text{HD-dist} \left(\xi_{n+j+1}(\omega), f_a^j \left(\xi_{n+1}(\omega) \right) \right) < \frac{1}{1000} \left| f_a^j \left(\xi_{n+1}(\omega) \right) \right|$

for all $j \leq p(\mu, \omega)$.

2.3. Escape-times. An important role is played by the *escape-times* of a . They are an infinite subsequence of $\{n_k(a)\}$, defined via the condition

$$(2.4) \quad \begin{aligned} & n_k \text{ is an escape-time} \\ & \iff \\ & \xi_{n_k}(\omega_{n_{k-1}}) \text{ intersects } (-\delta^2, \delta^2) \text{ and } |\xi_{n_k}(\omega_{n_{k-1}})| \geq \delta. \end{aligned}$$

It will be important that each $a \in \mathcal{A}$ experiences infinitely many escape-times. This is a consequence of the second parameter selection principle, which roughly speaking discards parameters that on the average has to wait too long for their escape-situations.

3. ESCAPING

We prepare the proofs of theorems A and B by showing that any escaping parameter interval $\omega_n(a)$ contains a super-stable parameter as well as parameters for which the critical orbit lands on any prescribed point in any hyperbolic set of f_a .

Lemma 5. *Pick an $a \in \mathcal{A}$ and a hyperbolic set Γ_a for f_a and a point $z(a) \in \Gamma_a$. Let n_k be an escape time for a . Then there are two parameters $a^*, \hat{a} \in \omega_{n_k-1}(a)$ and two integers r^* and $\hat{r}$, $0 < r^* \leq \hat{r} \lesssim -\log \delta$ such that*

- (i) f_{a^*} has a super-stable attractor of period $n_k(a) + r^*$;
- (ii) $f_{\hat{a}}^j(0) \neq z(\hat{a})$ for $j < n_k + \hat{r}$ and $f_{\hat{a}}^{n_k + \hat{r}}(0) = z(\hat{a})$.

Proof. Let $\omega_{n_k-1}(a) = (b, c)$. We may assume that $\xi_{n_k}(b) = \delta^2$ and $\xi_{n_k}(c) = \delta$. The idea is that distance between $\xi_{n_k+j}(b)$ and -1 will be $\leq 4^j \delta^4$ for $j > 1$, while the distance between -1 and $\xi_{n_k+j}(c)$ will be $\geq 3^j \delta^2$ as long as $\xi_{n_k+j}(c) < -3/4$. From this it follows that for some $j_0 \lesssim -\log \delta$, $\xi_{n_k+j_0}(\omega_{n_k})$ will grow to length $\sim 1/4$, with its left end $\xi_{n_k+j_0}(b)$ still within a distance $o(\delta)$ from -1 .

Thus $\xi_{n_k+r^*}$ maps (b, c) across $x = 0$ for $r^* = j_0 + 1$ or $r^* = j_0 + 2$, and since $0 \notin \xi_j(b, c)$ for $j < n_k + j^*$ by (2.2), the required a^* has been found.

Since Γ_a is hyperbolic, $z(a)$ moves continuously with a , and $(z(b), z(c))$ will be a very small interval. It follows that $\xi_{n_k+\hat{r}}(b, c) \supset (z(b), z(c))$ for $\hat{r} = r^*$ or $\hat{r} = r^* + 1$. Thus $\xi_{n_k+\hat{r}}(a) - z(a)$ changes sign on (b, c) , and so $\xi_{n_k+\hat{r}}(\hat{a}) = z(\hat{a})$ for some $\hat{a} \in (b, c)$. $\square$

4. PROOF OF THEOREM A AND THEOREM B

Each $a \in \mathcal{A}$ is given as

$$a = \cap_{n=1}^{\infty} \omega_n, \quad \omega_n \in \mathcal{P}_n,$$

with infinitely many escape situations where Lemma 5 can be applied. For any $b \in \omega_n$, inequality (2.1) holds. Thus we may use Lemma 1 to conclude that $|\omega_n| \lesssim e^{-n\lambda}$. From this the first two statements of Theorem A and Theorem B follows.

We now prove the last part of Theorem A. Let $a \in \mathcal{A}$ be such that $x = 0$ is generic for μ_a , and let $\{a_n\}$ be the sequence of super-stable parameters converging to a constructed above.

We have to show that

$$\lim_{n \rightarrow \infty} \int \varphi d\mu_{a_n} = \int \varphi d\mu_a$$

for any continuous function φ . It is enough to consider Lipschitz continuous test functions. Let κ be the Lipschitz' constant of φ , and let r_k denote the length of super-stable attractor of f_{a_k} . We have that for any a_k ,

$$\begin{aligned} \left| \int \varphi d\mu_a - \int \varphi d\mu_{a_k} \right| &\leq \left| \int \varphi d\mu_a - \frac{1}{r_k} \sum_{j=0}^{r_k-1} \varphi(\xi_j(a)) \right| \\ &\quad + \left| \frac{1}{r_k} \sum_{j=0}^{r_k-1} (\varphi(\xi_j(a)) - \varphi(\xi_j(a_k))) \right| \\ &\quad + \left| \frac{1}{r_k} \sum_{j=0}^{r_k-1} \varphi(\xi_j(a_k)) - \int \varphi d\mu_{a_k} \right|, \end{aligned}$$

where the last term = 0 by definition, and the first one is $< \epsilon$ for all $k > N_0(\epsilon)$ since 0 is generic for μ_a and since $r_k \uparrow \infty$ when $k \rightarrow \infty$. The second term finally is

$$\leq \frac{\kappa}{r_k} \sum_{j=0}^{r_k-1} |\xi_j(a) - \xi_j(a_k)|,$$

so it suffices to prove $S := \sum_{j=0}^{r_k-1} |\xi_j(a) - \xi_j(a_k)|$ is bounded by some constant independent of a_k . Let

$$\Lambda_j = |\xi_j(a) - \xi_j(a_k)|,$$

and let n_k be the escape-time preceding the creation of the super-stable orbit at time r_k ; then $r_k - n_k \lesssim \log 1/\delta$ (c.f. Lemma 5). Remember that $a_k \in \omega_{n_k}(a) \in \mathcal{P}_{n_k}$, and that for the “orbit” of ω_{n_k} under the family $\{\xi_i\}_{i=1}^{n_k}$ certain free return times $\{t_i\}_{i=1}^T$ are defined, $t_T = n_k$, and that each free return t_i is followed by a bound period of finite length, which we denote p_i . For the sake of notation, we define $t_0 = p_0 = 0$. We split the sum S into sub-sums:

$$S := \sum_{i=0}^{T-1} \left(S_i^{\text{bp}} + S_i^{\text{fp}} \right) + S^{\text{tail}},$$

where

$$S_i^{\text{bp}} = \sum_{l=t_i}^{t_i+p_i-1} \Lambda_l, \quad S_i^{\text{fp}} = \sum_{l=t_{i+1}}^{t_{i+1}-1} \Lambda_l$$

and

$$S^{\text{tail}} = \sum_{l=t_T}^{r_k} \Lambda_l.$$

With this notation, S_0^{fp} is the contribution up till the first free return and S_0^{bp} is an empty sum and equals 0. First we observe that S^{tail} has no more than $\sim -\log \delta$ terms, and is therefore $\leq C(\delta)$. We now estimate $\sum_{i=0}^{T-1} S_i^{\text{fp}}$, using (i) and (ii) of Lemma 3:

$$\begin{aligned} \sum_{i=0}^{T-1} S_i^{\text{fp}} &= \sum_{i=0}^{T-1} \sum_{l=t_i+p_i}^{t_{i+1}-1} \Lambda_l \leq \sum_{i=0}^{T-1} \sum_{l=t_i+p_i}^{t_{i+1}-1} C e^{-\lambda(t_{i+1}-l)} \Lambda_{t_{i+1}} \\ &\leq C_1 \sum_{i=1}^T \Lambda_{t_i} \leq C_1 \sum_{i=1}^T 2^{i-T} \Lambda_{t_T} \leq C_2. \end{aligned}$$

We now turn to $\sum_{i=0}^{T-1} S_i^{\text{bp}}$. First we estimate the individual terms in S_i^{bp} . Now $\Lambda_l = |\xi_l(a) - \xi_l(a_k)|$, and $(a; a_k) \subset \omega_{n_k} \in \mathcal{P}_{n_k}$. Since t_i is a free return, $t_i < t_T = n_k$, it follows that $\xi_{t_i}((a; a_k)) \subset \text{some } I_{\mu_i \nu_i}^+ \subset I^*$. Using Lemma 2, Lemma 4 and the binding condition (2.3), we see that for $1 \leq j < p_i$ and any $b \in (a; a_k)$,

$$\begin{aligned} \Lambda_{t_i+j} &\lesssim |f_b^j(\xi_{t_i}((a; a_k)))| = \frac{|f_b^j(\xi_{t_i}((a; a_k)))|}{|f_b^j(\hat{I}_{\mu_i})|} |f_b^j(\hat{I}_{\mu_i})| \\ &\lesssim \frac{|f_b(\xi_{t_i}((a; a_k)))|}{|f_b(\hat{I}_{\mu_i})|} e^{-2\alpha j} \sim \frac{|f_b(\xi_{t_i}((a; a_k)))|}{|f_b(I_{\mu_i})|} e^{-2\alpha j} \\ &\sim \frac{|(\xi_{t_i}((a; a_k)))|}{|I_{\mu_i}|} e^{-2\alpha j} \end{aligned}$$

In the last two steps we also used the facts that $|f_b(\hat{I}_{\mu})| \sim |f_b(I_{\mu})|$, and that the distortion of f_b restricted to I_{μ} has a bound independent of μ .

Obviously $\Lambda_{t_j} < |\xi_{t_i}((a; a_k))|/|I_{\mu_i}|$, so we obtain

$$S_i^{\text{bp}} \lesssim \sum_{j=0}^{\infty} \frac{|(\xi_{t_i}((a; a_k)))|}{|I_{\mu_i}|} e^{-2\alpha j} \lesssim \frac{|(\xi_{t_i}((a; a_k)))|}{|I_{\mu_i}|} \leq 3 \frac{|I_{\mu_i \nu_i}|}{|I_{\mu_i}|} \lesssim \frac{1}{\mu_i^2}.$$

Finally we estimate $\sum S_i^{\text{bp}}$:

$$\begin{aligned} \sum_{i=0}^{T-1} S_i^{\text{bp}} &\lesssim \sum_{\mu_i} \sum_{\substack{\text{returns} \\ \text{to } I_{\mu_i}}} \mu_i^{-2} \lesssim \sum_{\mu_i} \sum_{\substack{\text{last return} \\ \text{to } I_{\mu_i}}} \mu_i^{-2} \\ &\lesssim \sum_{|\mu| \geq -\log \delta} \mu^{-2} < C. \end{aligned}$$

The second “ $\lesssim$ ” holds because of (ii) of Lemma 3. This finishes the proof of Theorem A.

5. PROOF OF THEOREM D

Suppose $\Gamma_a = \{x_1, x_2, \dots, x_p\}$ is a hyperbolic repelling periodic orbit for f_a , absorbing the critical orbit. Let N be minimal such that $f_a^N(0) \in \Gamma_a$, and assume that $f_a^N(0) = x_1$. For $\gamma > 0$ we define $J_\gamma = \bigcup_{i=1}^p [x_i - \gamma, x_i + \gamma]$.

Since f_a is Misiurewicz map with finite critical orbit, we can carry out the Benedicks-Carleson construction in a neighborhood of a and find a sequence $\{\omega_n\}_{n=N}^\infty$ of parameter intervals such that

- $\omega_{n+1} \subset \omega_n$ and $a = \bigcap_{n=N}^\infty \omega_n$;
- $\xi_k(\omega_n) \subset J_\gamma$ for $k = N, \dots, n$;
- for all $n \geq N$ there is an i_n such that $\xi_n(\omega_n) = [x_{i_n} - \gamma, x_{i_n} + \gamma]$;
- there is a natural number $\hat{N} = \hat{N}(\gamma)$ such that for each $n \geq N$ there is an $m_n \leq \hat{N}$ such that $\xi_{n+m_n}(\omega_n) \ni 0$.

This follows from continuity, uniform expansion away from the critical point and the fact that parameter- and space-derivatives are comparable. It follows that for all $n \geq N$, there is an $a_n \in \omega_n$ such that f_{a_n} has super-attractor of length $n + m_n$ such that

$$\#\{j \leq n + m_n \mid f_{a_n}^j(0) \notin J_\gamma\} = N - 1 + m_n \leq N + \hat{N}(\gamma).$$

Now pick a $\varphi \in C^0(I)$ and an $\epsilon > 0$. Since φ is uniformly continuous, we can fix a γ such that $|x - y| < \gamma$ implies $|\varphi(x) - \varphi(y)| < \epsilon/2$. If μ_{a_n} is the natural measure for f_{a_n} , where a_n is as above, we have that

$$\begin{aligned} \int \varphi d\mu_{a_n} &= \frac{1}{n + m_n} \sum_{i=1}^{n+m_n} \varphi(f_{a_n}^i(0)) \\ &\leq \frac{1}{n + m_n} \sum_{i=N}^n \varphi(f_{a_n}^i(0)) + \frac{N - 1 + m_n}{n + m_n} \sup_I \varphi \\ &\leq \frac{1}{n + m_n} \sum_{i=N}^n \varphi(f_{a_n}^i(0)) + \frac{N + \hat{N}}{n + \hat{N}} \sup_I \varphi. \end{aligned}$$

It is clear that the last term vanishes when $n \rightarrow \infty$. We have to show that $\frac{1}{n+m_n} \sum_{i=N}^n \varphi(f_{a_n}^i(0))$ tends to $\int \varphi d\mu_a^{\text{sing}} = \frac{1}{p} \sum_{i=1}^p \varphi(x_i)$, when n tends to infinity. Without loss of generality we may assume that $n - N + 1 = R_n p$ for some integer R_n . Let $y_i = f_{a_n}^{N+i-1}(0)$ for $1 \leq i \leq n + 1 - N$, and for $i > p$ define $x_i = f_a^{i-1}(x_1)$. Then

$$\begin{aligned} & \frac{1}{n+m_n} \sum_{i=N}^n \varphi(f_{a_n}^i(0)) \\ &= \frac{1}{n+m_n} \sum_{i=1}^{n+1-N} \varphi(y_i) \\ &\leq \frac{pR_n}{n+m_n} \frac{1}{pR_n} \sum_{i=1}^{n+1-N} (\varphi(x_i) + \epsilon/2) \\ &= \frac{n-N+1}{n+m_n} \left(\int \varphi d\mu_a^{\text{sing}} + \epsilon/2 \right) \\ &\leq \int \varphi d\mu_a^{\text{sing}} + \epsilon, \end{aligned}$$

where the last inequality holds for n sufficiently large. In the same way

$$\int \varphi d\mu_{a_n} \geq \int \varphi d\mu_a^{\text{sing}} - \epsilon$$

for n sufficiently large.

6. PROOF OF THEOREM C

Since $\{x_1, \dots, x_p\}$ is a hyperbolic repeller, it persists for nearby parameter values. Let $\Gamma_a = \{x_1(a), \dots, x_p(a)\}$ by its continuation. By Theorem B, we know that there is a sequence $\{b_n\}$ converging to a , such that f_{b_n} has critical point pre-periodic to $x_1(b_n)$. By Theorem D, for each b_n there is a sequence $\{a_{n,m}\}_{m=1}^\infty$ corresponding to maps with super stable attractors, such that $\lim_{m \rightarrow \infty} a_{n,m} = b_n$ and $\lim_{m \rightarrow \infty} \mu_{a_{n,m}} = \sum_{i=1}^p \delta_{x_i(b_n)}$. Since Γ_a moves continuously with a , we get a sequence $\{a_{n,m(n)}\}_{n=1}^\infty$, corresponding to super attractors and converging to a such that

$$\lim_{n \rightarrow \infty} \mu_{a_{n,m(n)}} \stackrel{\text{weak}^*}{=} \sum_{i=1}^p \delta_{x_i(a)}.$$

REFERENCES

- [BC85] M. Benedicks and L. Carleson, *On iterations of $1 - ax^2$* , Ann. of Math. **122** (1985), 1–25.

- [BC91] M. Benedicks and L. Carleson, *The dynamics of the Hénon map*, Ann. of Math. **133** (1991), 73–169.
- [BL91] A.M. Blokh and M.Yu. Lyubich, *Measurable dynamics of S -unimodal maps of the interval*, Ann. Sc. E.N.S. 4e série **24** (1991), 545–573.
- [BM89] M. Benedicks and M. Misiurewicz, *Absolutely continuous invariant measures for maps with flat tops*, Inst. Hautes Études Sci. Publ. Math. (1989), no. 69, 203–213.
- [BV96] V. Baladi and M. Viana, *Strong stochastic stability and rate of mixing for unimodal maps*, Ann. Sci. cole Norm. Sup. (4) **29** (1996), no. 4, 483–517.
- [BY92] M. Benedicks and L. S. Young, *Absolutely continuous invariant measures and random perturbations for certain one-dimensional maps*, Ergod. Th. & Dynam. Sys. **12** (1992), 13–37.
- [HK90] F. Hofbauer and G. Keller, *Quadratic maps without asymptotic measure*, Comm. Math. Phys. **127** (1990), no. 2, 319–337.
- [Jak81] M. Jakobson, *Absolutely continuous invariant measures for one-parameter families of one-dimensional maps*, Comm. Math. Phys. **81** (1981), 39–88.
- [Joh87] S.D. Johnson, *Singular measures without restrictive intervals*, Comm. Math. Phys. **110** (1987), no. 2, 185–190.
- [Luz98] S. Luzzatto, *A proof of Jakobson’s theorem*, In preparation, 1998.
- [Lyu97] M. Lyubich, *Almost every real quadratic map is either regular or stochastic*, SUNY preprint IMS 97–8, SUNY Stony-Brook, New York, 1997.
- [Mis81] M. Misiurewicz, *Absolutely continuous measures for certain maps of an interval*, Inst. Hautes Études Sci. Publ. Math. (1981), no. 53, 17–51.
- [MS93] W. de Melo and S. van Strien, *One-dimensional dynamics*, Springer-Verlag, Berlin-Heidelberg, 1993.
- [RS92] M. Rychlik and E. Sorets, *Regularity and other properties of absolutely continuous invariant measures for the quadratic family*, Comm. Math. Phys. **150** (1992), no. 2, 217–236.
- [Ryc88] M. Rychlik, *Another proof of Jakobson’s theorem and related results*, Ergod. Th. & Dynam. Sys. **8** (1988), 93–109.
- [Thu96] H. Thunberg, *Absolutely continuous invariant measures and super-stable periodic orbits: weak*-convergence of natural measures*, Some Problems in Unimodal Dynamics (Stockholm), KTH, Stockholm, 1996, Ph.D. thesis, pp. I: 1–19.
- [Thu97] H. Thunberg, *Positive exponent in families with flat critical point*, To appear in Ergod. Th. and Dynam. Sys., 1997.
- [Tsu93a] M. Tsujii, *Positive Lyapunov exponent in families of one-dimensional dynamical systems*, Inven. math. **111** (1993), no. 1, 113–137.
- [Tsu93b] M. Tsujii, *A proof of Benedicks-Carleson-Yakobson theorem*, Tokyo J. Math. **16** (1993), no. 2, 295–310.
- [Tsu96] M. Tsujii, *On continuity of Bowen-Ruelle-Sinai measures in families of one-dimensional maps*, Comm. Math. Phys. **117** (1996), no. 1, 1–11.
- [TTY92] P. Thieullen, C. Tresser, and L.S. Young, *Exposant de Lyapunov positif dans des familles à un paramètre d’applications unimodales*, C. R. Acad. Sci. Paris, Série I **t315** (1992), 69–72.
- [Ure95] R. Ures, *On the approximation of Hénon-like attractors by homoclinic tangencies*, Ergodic Theory Dynam. Systems **15** (1995), no. 6, 1223–1229.

- [Ure96] R. Ures, *Hénon attractors: SBR measures and Dirac measures for sinks*, International Conference on Dynamical Systems (Montevideo, 1995) (Harlow), Pitman Res. Notes Math. Ser., vol. 362, Longman, Harlow, 1996, pp. 214–219.

DEPARTMENT OF MATHEMATICS, KTH, S-100 44 STOCKHOLM, SWEDEN
E-mail address: `hasset@math.kth.se`